







INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C08B 11/20, 15/00, C11D 3/22	A1	(11) International Publication Number: WO 99/14245 (43) International Publication Date: 25 March 1999 (25.03.99)
<p>(21) International Application Number: PCT/US98/19139</p> <p>(22) International Filing Date: 15 September 1998 (15.09.98)</p> <p>(30) Priority Data: 60/058,892 15 September 1997 (15.09.97) US</p> <p>(71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): LEUPIN, Jennifer, Ann [CA/US]; 7974 Fawncreek Drive, Cincinnati, OH 45249 (US). BOYER, Stanton, Lane [US/US]; 12 Fall River Court, Fairfield, OH 45014 (US). GOSSELINK, Eugene, Paul [US/US]; 3754 Susanna Drive, Cincinnati, OH 45251 (US). WANG, Jiping [CN/US]; 6546 Glenarbor Drive, West Chester, OH 45069 (US). HUNTER, Kathleen, Brenner [US/US]; 855 Rogers Road, Villa Hills, KY 41017 (US). WASHINGTON, Nodie [US/US]; 1089 Wilberforce-Clifton Road, Wilberforce, OH 45384 (US).</p> <p>(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).</p>	<p>(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: **LAUNDRY DETERGENT COMPOSITIONS WITH CELLULOSIC BASED POLYMERS TO PROVIDE APPEARANCE AND INTEGRITY BENEFITS TO FABRICS LAUNDERED THEREWITH**

(57) Abstract

Compositions and methods that utilize certain cellulosic based polymer or oligomer materials as fabric treatment agents which can impart fabric appearance and integrity benefits to fabrics and textiles laundered in washing solutions containing such materials. The cellulosic based polymer or oligomer materials can be characterized by general formula: (I) wherein each R is independently selected from the group consisting of R_C, R_H, and (II). Each R₂ is independently selected from the group consisting of H and C₁-C₄ alkyl. Each R_C is (III); and each R_H is (IV). The Degree of Substitution for group R_H is between about 0.001 and 0.3, and the Degree of Substitution for group R_C wherein Z is H or M is between about 0.2 and 2.0.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**LAUNDRY DETERGENT COMPOSITIONS WITH CELLULOSIC BASED POLYMERS TO
PROVIDE APPEARANCE AND INTEGRITY BENEFITS TO FABRICS LAUNDERED
THEREWITH**

TECHNICAL FIELD

The present invention relates to compositions, in either liquid or granular form, for use in laundry applications, wherein the compositions comprise certain cellulosic based polymer or oligomer materials which impart appearance and integrity benefits to fabrics and textiles laundered in washing solutions formed from such compositions.

BACKGROUND OF THE INVENTION


It is, of course, well known that alternating cycles of using and laundering fabrics and textiles, such as articles of worn clothing and apparel, will inevitably adversely affect the appearance and integrity of the fabric and textile items so used and laundered. Fabrics and textiles simply wear out over time and with use. Laundering of fabrics and textiles is necessary to remove soils and stains which accumulate therein and thereon during ordinary use. However, the laundering operation itself, over many cycles, can accentuate and contribute to the deterioration of the integrity and the appearance of such fabrics and textiles.

Deterioration of fabric integrity and appearance can manifest itself in several ways. Short fibers are dislodged from woven and knit fabric/textile structures by the mechanical action of laundering. These dislodged fibers may form lint, fuzz or "pills" which are visible on the surface of fabrics and diminish the appearance of newness of the fabric. Further, repeated laundering of fabrics and textiles, especially with bleach-containing laundry products, can remove dye from fabrics and textiles and impart a faded, worn out appearance as a result of diminished color intensity, and in many cases, as a result of changes in hues or shades of color.


Given the foregoing, there is clearly an ongoing need to identify materials which could be added to laundry detergent products that would associate themselves with the fibers of the fabrics and textiles laundered using such detergent products and thereby reduce or minimize the tendency of the laundered fabric/textiles to deteriorate in appearance. Any such detergent product additive material should, of course, be able to benefit fabric appearance and integrity without unduly interfering with the ability of the laundry detergent to perform its fabric cleaning function. The present invention is directed to the use of cellulosic based polymer or oligomer materials in laundry applications which perform in this desired manner.

SUMMARY OF THE INVENTION

Cellulosic based polymer or oligomer materials which are suitable for use in laundry operations and provide the desired fabric appearance and integrity benefits can be characterized by the following general formula:




wherein each R is independently selected from the group consisting of R_C , R_H , and




wherein:

- each R_2 is independently selected from the group consisting of H and C_1 - C_4 alkyl;
- each R_C is



- each R_H is



wherein:

- each Z is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl, hydroxyalkyl, and M;
- each R_5 is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl and hydroxyalkyl

wherein:

M is selected from the group consisting of Na, K, $1/2\text{Ca}$, and $1/2\text{Mg}$;

each x is from 0 to about 5;

each y is from about 1 to about 5;

provided that:

-if Z bears a positive charge, it is balanced by a suitable anion; and

-the Degree of Substitution for group R_H is between about 0.001 and 0.3, more preferably between about 0.005 and 0.2, and most preferably between about 0.01 and 0.1; and

-the Degree of Substitution for group R_C wherein Z is H or M is between about 0.2 and 2.0, more preferably between about 0.3 and 1.0, and most preferably between about 0.4 and 0.7.

In another embodiment of the present invention a process is defined wherein cellulose derivatives are obtained by conversion of a carboxylalkylated cellulosic at least partially to an ester, followed by treatment with an amine in the presence of at least traces of water. Alternatively, the cellulose derivatives obtained by conversion of a carboxylalkylated cellulosic at least partially to an ester, followed by treatment with enough base in the presence of water to hydrolyze at least 10% of the ester groups.

The cellulosic based polymer or oligomer materials defined above can be used as a washing solution additive in either granular or liquid form. Alternatively, they can be admixed to granular detergents, dissolved in liquid detergent compositions or added to a fabric softening composition. The forgoing description of uses for the cellulosic based fabric treatment materials defined herein are intended to be exemplary and other uses will be apparent to those skilled in the art and are intended to be within the scope of the present invention.

The laundry detergent compositions herein comprise from about 1% to 80% by weight of a deterative surfactant, from about 0.1% to 80% by weight of an organic or inorganic detergency builder and from about 0.1% to 5% by weight of the cellulosic based fabric treatment materials of the present invention. The deterative surfactant and detergency builder materials can be any of those useful in conventional laundry detergent products.

Aqueous solutions of the cellulosic based polymer or oligomer materials of the subject invention comprise from about 0.1% to 80% by weight of the cellulosic based fabric treatment materials dissolved in water and other ingredients such as stabilizers and pH adjusters.

In its method aspect, the present invention relates to the laundering or treating of fabrics and textiles in aqueous washing or treating solutions formed from effective amounts of the detergent compositions described herein, or formed from the individual components of such compositions.

Laundrying of fabrics and textiles in such washing solutions, followed by rinsing and drying, imparts fabric appearance benefits to the fabric and textile articles so treated. Such benefits can include improved overall appearance, pill/fuzz reduction, antifading, improved abrasion resistance, and/or enhanced softness.

DETAILED DESCRIPTION OF THE INVENTION

As noted, when fabric or textiles are laundered in wash solutions which comprise the cellulosic based polymer or oligomer materials of the present invention fabric appearance and integrity are enhanced. The cellulosic based fabric treatment materials can be added to wash solutions by incorporating them into a detergent composition, a fabric softener or by adding them separately to the washing solution. The cellulosic based fabric treatment materials are described herein primarily as liquid or granular detergent additives but the present invention is not meant to be so limited. The cellulosic based fabric treatment materials, detergent composition components, optional ingredients for such compositions and methods of using such compositions, are described in detail below. All percentages are by weight unless other specified.


A) Cellulosic Based Polymer or Oligomer Materials

The essential component of the compositions of the present invention comprises one or more cellulosic based polymer or oligomer. Such materials have been found to impart a number of appearance benefits to fabrics and textiles laundered in aqueous washing solutions formed from detergent compositions which contain such cellulosic based fabric treatment materials. Such fabric appearance benefits can include, for example, improved overall appearance of the laundered fabrics, reduction of the formation of pills and fuzz, protection against color fading, improved abrasion resistance, etc. The cellulosic based fabric treatment materials used in the compositions and methods herein can provide such fabric appearance benefits with acceptably little or no loss in cleaning performance provided by the laundry detergent compositions into which such materials are incorporated.


As will be apparent to those skilled in the art, an oligomer is a molecule consisting of only a few monomer units while polymers comprise considerably more monomer units. For the present invention, oligomers are defined as molecules having an average molecular weight below about 1,000 and polymers are molecules having an average molecular weight of greater than about 1,000. One suitable type of cellulosic based polymer or oligomer fabric treatment material for use herein has an average molecular weight of from about 10,000 to about 2,000,000, preferably from about 50,000 to about 1,000,000.

The cellulosic based fabric treatment component of the detergent compositions herein will generally comprise from about 0.1% to about 5% by the weight of the detergent composition. More preferably, such cellulosic based fabric treatment materials will comprise from about 0.5% to about 4% by weight of the detergent compositions, most preferably from about 0.75% to about 3%. However, as discussed above, when used as a washing solution additive, i.e. when the cellulosic based fabric treatment component is not incorporated into a detergent composition, the concentration of the cellulosic based component can comprise from about 0.1% to about 80% by weight of the additive material.

One suitable group of cellulosic based polymer or oligomer materials for use herein is characterized by the following formula:




wherein each R is independently selected from the group consisting of R_C , R_H , and




wherein:

- each R_2 is independently selected from the group consisting of H and C_1 - C_4 alkyl;
- each R_C is



- each R_H is



wherein:

- each Z is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl,

alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl, hydroxyalkyl, and M;

- each R_5 is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl and hydroxyalkyl

wherein:

M is selected from the group consisting of Na, K, $1/2Ca$, and $1/2Mg$;

each x is from 0 to about 5;

each y is from about 1 to about 5;

provided that:

-if Z bears a positive charge, it is balanced by a suitable anion; and

-the Degree of Substitution for group R_H is between about 0.001 and 0.3, more preferably between about 0.005 and 0.2, and most preferably between about 0.01 and 0.1; and

-the Degree of Substitution for group R_C wherein Z is H or M is between about 0.2 and 2.0, more preferably between about 0.3 and 1.0, and most preferably between about 0.4 and 0.7.

The "Degree of Substitution" for group R_H , which is sometimes abbreviated herein " DS_{RH} ", means the number of moles of group R_H components that are substituted per anhydrous glucose unit, wherein an anhydrous glucose unit is a six membered ring as shown in the repeating unit of the general structure above.

The "Degree of Substitution" for group R_C , which is sometimes abbreviated herein " DS_{RC} ", means the number of moles of group R_C components, wherein Z is H or M, that are substituted per anhydrous glucose unit, wherein an anhydrous glucose unit is a six membered ring as shown in the repeating unit of the general structure above. The requirement that Z be H or M is necessary to insure that there are a sufficient number of carboxy methyl groups such that the resulting polymer is soluble. It is understood that in addition to the required number of R_C components wherein Z is H or M, there can be, and most preferably are, additional R_C components wherein Z is a group other than H or M.

In an especially preferred embodiment of this invention at least 0.1 of the R_3 groups are derived from $-CH_2-COOH$ or a salt thereof, that is the degree of substitution in the cellulose chain is at least 0.1 for CH_2-COOH and salts thereof. In another embodiment of the present invention all of the x's are equal to 0.

The subject cellulosic based polymer or oligomer materials can be produced by conversion of a carboxylalkylated cellulosic at least partially to an ester, followed by treatment with an amine in the presence of at least traces of water. Alternatively, the cellulose derivatives obtained by conversion of a carboxylalkylated cellulosic at least partially to an ester, followed by treatment with enough base in the presence of water to hydrolyze at least 10% of the ester groups. The production of materials according to the present invention is further defined in the Examples below.

B) Deterstive Surfactant

The detergent compositions herein comprise from about 1% to 80% by weight of a deterstive surfactant. Preferably such compositions comprise from about 5% to 50% by weight of surfactant. Deterstive surfactants utilized can be of the anionic, nonionic, zwitterionic, ampholytic or cationic type or can comprise compatible mixtures of these types. Detergent surfactants useful herein are described in U.S. Patent 3,664,961, Norris, issued May 23, 1972, U.S. Patent 3,919,678, Laughlin et al., issued December 30, 1975, U.S. Patent 4,222,905, Cockrell, issued September 16, 1980, and in U.S. Patent 4,239,659, Murphy, issued December 16, 1980. All of these patents are incorporated herein by reference. Of all the surfactants, anionics and nonionics are preferred.


Useful anionic surfactants can themselves be of several different types. For example, water-soluble salts of the higher fatty acids, i.e., "soaps", are useful anionic surfactants in the compositions herein. This includes alkali metal soaps such as the sodium, potassium, ammonium, and alkylammonium salts of higher fatty acids containing from about 8 to about 24 carbon atoms, and preferably from about 12 to about 18 carbon atoms. Soaps can be made by direct saponification of fats and oils or by the neutralization of free fatty acids. Particularly useful are the sodium and potassium salts of the mixtures of fatty acids derived from coconut oil and tallow, i.e., sodium or potassium tallow and coconut soap.

Additional non-soap anionic surfactants which are suitable for use herein include the water-soluble salts, preferably the alkali metal, and ammonium salts, of organic sulfuric reaction products having in their molecular structure an alkyl group containing from about 10 to about 20 carbon atoms and a sulfonic acid or sulfuric acid ester group. (Included in the term "alkyl" is the alkyl portion of acyl groups.) Examples of this group of synthetic surfactants are a) the sodium, potassium and ammonium alkyl sulfates, especially those obtained by sulfating the higher alcohols (C₈-C₁₈ carbon atoms) such as those produced by reducing the glycerides of tallow or coconut oil; b) the sodium, potassium and ammonium alkyl polyethoxylate sulfates, particularly those in which the alkyl group contains from 10 to 22, preferably from 12 to 18 carbon atoms, and wherein the polyethoxylate chain contains from 1 to 15, preferably 1 to 6 ethoxylate moieties; and c) the

sodium and potassium alkylbenzene sulfonates in which the alkyl group contains from about 9 to about 15 carbon atoms, in straight chain or branched chain configuration, e.g., those of the type described in U.S. Patents 2,220,099 and 2,477,383. Especially valuable are linear straight chain alkylbenzene sulfonates in which the average number of carbon atoms in the alkyl group is from about 11 to 13, abbreviated as C₁₁₋₁₃ LAS.


Preferred nonionic surfactants are those of the formula R₁(OC₂H₄)_nOH, wherein R₁ is a C₁₀-C₁₆ alkyl group or a C₈-C₁₂ alkyl phenyl group, and n is from 3 to about 80. Particularly preferred are condensation products of C₁₂-C₁₅ alcohols with from about 5 to about 20 moles of ethylene oxide per mole of alcohol, e.g., C₁₂-C₁₃ alcohol condensed with about 6.5 moles of ethylene oxide per mole of alcohol.

Additional suitable nonionic surfactants include polyhydroxy fatty acid amides of the formula:






wherein R is a C₉₋₁₇ alkyl or alkenyl, R₁ is a methyl group and Z is glycityl derived from a reduced sugar or alkoxyated derivative thereof. Examples are N-methyl N-1-deoxyglucityl cocoamide and N-methyl N-1-deoxyglucityl oleamide. Processes for making polyhydroxy fatty acid amides are known and can be found in Wilson, U.S. Patent 2,965,576 and Schwartz, U.S. Patent 2,703,798, the disclosures of which are incorporated herein by reference.

Preferred surfactants for use in the detergent compositions described herein are amine based surfactants of the general formula:




wherein R₁ is a C₆-C₁₂ alkyl group; n is from about 2 to about 4, X is a bridging group which is selected from NH, CONH, COO, or O or X can be absent; and R₃ and R₄ are individually selected from H, C₁-C₄ alkyl, or (CH₂-CH₂-O(R₅)) wherein R₅ is H or methyl. Especially preferred amines based surfactants include the following:





wherein R_1 is a $\text{C}_6\text{-C}_{12}$ alkyl group and R_5 is H or CH_3 . Particularly preferred amines for use in the surfactants defined above include those selected from the group consisting of octyl amine, hexyl amine, decyl amine, dodecyl amine, $\text{C}_8\text{-C}_{12}$ bis(hydroxyethyl)amine, $\text{C}_8\text{-C}_{12}$ bis(hydroxyisopropyl)amine, $\text{C}_8\text{-C}_{12}$ amido-propyl dimethyl amine, or mixtures thereof.

In a highly preferred embodiment, the amine based surfactant is described by the formula:



wherein R_1 is $\text{C}_8\text{-C}_{12}$ alkyl.

C) Detergent Builder

The detergent compositions herein may also comprise from about 0.1% to 80% by weight of a detergent builder. Preferably such compositions in liquid form will comprise from about 1% to 10% by weight of the builder component. Preferably such compositions in granular form will comprise from about 1% to 50% by weight of the builder component. Detergent builders are well known in the art and can comprise, for example, phosphate salts as well as various organic and inorganic nonphosphorus builders.

Water-soluble, nonphosphorus organic builders useful herein include the various alkali metal, ammonium and substituted ammonium polyacetates, carboxylates, polycarboxylates and polyhydroxy sulfonates. Examples of polyacetate and polycarboxylate builders are the sodium, potassium, lithium, ammonium and substituted ammonium salts of ethylene diamine tetraacetic acid, nitrilotriacetic acid, oxydisuccinic acid, mellitic acid, benzene polycarboxylic acids, and citric acid. Other suitable polycarboxylates for use herein are the polyacetal carboxylates described in U.S. Patent 4,144,226, issued March 13, 1979 to Crutchfield et al., and U.S. Patent 4,246,495, issued March 27, 1979 to Crutchfield et al., both of which are incorporated herein by reference. Particularly preferred polycarboxylate builders are the oxydisuccinates and the ether carboxylate builder compositions comprising a combination of tartrate monosuccinate and tartrate disuccinate described in U.S. Patent 4,663,071, Bush et al., issued May 5, 1987, the disclosure of which is incorporated herein by reference.

Examples of suitable nonphosphorus, inorganic builders include the silicates, aluminosilicates, borates and carbonates. Particularly preferred are sodium and potassium carbonate, bicarbonate, sesquicarbonate, tetraborate decahydrate, and silicates having a weight ratio of SiO_2 to alkali metal oxide of from about 0.5 to about 4.0, preferably from about 1.0 to

about 2.4. Also preferred are aluminosilicates including zeolites. Such materials and their use as detergent builders are more fully discussed in Corkill et al., U. S. Patent No. 4,605,509, the disclosure of which is incorporated herein by reference. Also discussed in U. S. Patent No. 4,605,509 are crystalline layered silicates which are suitable for use in the detergent compositions of this invention.

D) Optional Detergent Ingredients

In addition to the surfactants, builders and cellulosic based polymer or oligomer materials hereinbefore described, the detergent compositions of the present invention can also include any number of additional optional ingredients. These include conventional detergent composition components such as enzymes and enzyme stabilizing agents, suds boosters or suds suppressers, anti-tarnish and anticorrosion agents, soil suspending agents, soil release agents, germicides, pH adjusting agents, non-builder alkalinity sources, chelating agents, organic and inorganic fillers, solvents, hydrotropes, optical brighteners, dyes and perfumes.

pH adjusting agents may be necessary in certain applications where the pH of the wash solution is greater than about 10.0 because the fabric integrity benefits of the defined compositions begin to diminish at a higher pH. Hence, if the wash solution is greater than about 10.0 after the addition of the cellulosic based polymer or oligomer materials of the present invention a pH adjuster should be used to reduce the pH of the washing solution to below about 10.0, preferably to a pH of below about 9.5 and most preferably below about 7.5. Suitable pH adjusters will be known to those skilled in the art.


A preferred optional ingredients for incorporation into the detergent compositions herein comprises a bleaching agent, e.g., a peroxygen bleach. Such peroxygen bleaching agents may be organic or inorganic in nature. Inorganic peroxygen bleaching agents are frequently utilized in combination with a bleach activator.

Useful organic peroxygen bleaching agents include percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of metachloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in U.S. Patent 4,483,781, Hartman, Issued November 20, 1984; European Patent Application EP-A-133,354, Banks et al., Published February 20, 1985; and U.S. Patent 4,412,934, Chung et al., Issued November 1, 1983. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxy caproic acid (NAPAA) as described in U.S. Patent 4,634,551, Issued January 6, 1987 to Burns et al.

Inorganic peroxygen bleaching agents may also be used, generally in particulate form, in the detergent compositions herein. Inorganic bleaching agents are in fact preferred. Such inorganic peroxygen compounds include alkali metal perborate and percarbonate materials. For example, sodium perborate (e.g. mono- or tetra-hydrate) can be used. Suitable inorganic bleaching agents can also include sodium or potassium carbonate peroxyhydrate and equivalent "percarbonate" bleaches, sodium pyrophosphate peroxyhydrate, urea peroxyhydrate, and sodium peroxide. Persulfate bleach (e.g., OXONE, manufactured commercially by DuPont) can also be used. Frequently inorganic peroxygen bleaches will be coated with silicate, borate, sulfate or water-soluble surfactants. For example, coated percarbonate particles are available from various commercial sources such as FMC, Solvay Interlox, Tokai Denka and Degussa.

Inorganic peroxygen bleaching agents, e.g., the perborates, the percarbonates, etc., are preferably combined with bleach activators, which lead to the *in situ* production in aqueous solution (i.e., during use of the compositions herein for fabric laundering/bleaching) of the peroxy acid corresponding to the bleach activator. Various non-limiting examples of activators are disclosed in U.S. Patent 4,915,854, Issued April 10, 1990 to Mao et al.; and U.S. Patent 4,412,934 Issued November 1, 1983 to Chung et al. The nonanoyloxybenzene sulfonate (NOBS) and tetraacetyl ethylene diamine (TAED) activators are typical and preferred. Mixtures thereof can also be used. See also the hereinbefore referenced U.S. 4,634,551 for other typical bleaches and activators useful herein.


Other useful amido-derived bleach activators are those of the formulae:




wherein R^1 is an alkyl group containing from about 6 to about 12 carbon atoms, R^2 is an alkylene containing from 1 to about 6 carbon atoms, R^5 is H or alkyl, aryl, or alkaryl containing from about 1 to about 10 carbon atoms, and L is any suitable leaving group. A leaving group is any group that is displaced from the bleach activator as a consequence of the nucleophilic attack on the bleach activator by the perhydrolysis anion. A preferred leaving group is phenol sulfonate.

Preferred examples of bleach activators of the above formulae include (6-octanamido-caproyl)oxybenzenesulfonate, (6-nonanamidocaproyl) oxybenzenesulfonate, (6-decanamidocaproyl)oxybenzenesulfonate and mixtures thereof as described in the hereinbefore referenced U.S. Patent 4,634,551.

Another class of useful bleach activators comprises the benzoxazin-type activators disclosed by Hodge et al. in U.S. Patent 4,966, 723, Issued October 30, 1990, incorporated herein by reference. A highly preferred activator of the benzoxazin-type is:



Still another class of useful bleach activators includes the acyl lactam activators, especially acyl caprolactams and acyl valerolactams of the formulae:



wherein R^6 is H or an alkyl, aryl, alkoxyaryl, or alkaryl group containing from 1 to about 12 carbon atoms. Highly preferred lactam activators include benzoyl caprolactam, octanoyl caprolactam, 3,5,5-trimethylhexanoyl caprolactam, nonanoyl caprolactam, decanoyl caprolactam, undecenoyl caprolactam, benzoyl valerolactam, octanoyl valerolactam, nonanoyl valerolactam, decanoyl valerolactam, undecenoyl valerolactam, 3,5,5-trimethylhexanoyl valerolactam and mixtures thereof. See also U.S. Patent 4,545,784, Issued to Sanderson, October 8, 1985, incorporated herein by reference, which discloses acyl caprolactams, including benzoyl caprolactam, adsorbed into sodium perborate.

If utilized, peroxygen bleaching agent will generally comprise from about 2% to 30% by weight of the detergent compositions herein. More preferably, peroxygen bleaching agent will comprise from about 2% to 20% by weight of the compositions. Most preferably, peroxygen bleaching agent will be present to the extent of from about 3% to 15% by weight of the compositions herein. If utilized, bleach activators can comprise from about 2% to 10% by weight of the detergent compositions herein. Frequently, activators are employed such that the molar ratio of bleaching agent to activator ranges from about 1:1 to 10:1, more preferably from about 1.5:1 to 5:1.

Another highly preferred optional ingredient in the detergent compositions herein is a deterative enzyme component. While it is known that some enzymes will degrade the peptide bonds of cellulotics, the cellulosic based polymer or oligomer materials defined herein do not exhibit such degradation in the presence of enzymes. Hence, enzymes can be added to detergent compositions which comprise the cellulosic based fabric treatment materials of the present invention with substantially no degradation.

Enzymes can be included in the present detergent compositions for a variety of purposes, including removal of protein-based, carbohydrate-based, or triglyceride-based stains from substrates, for the prevention of refugee dye transfer in fabric laundering, and for fabric restoration. Suitable enzymes include proteases, amylases, lipases, cellulases, peroxidases, and mixtures thereof of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin. Preferred selections are influenced by factors such as pH-activity and/or stability, optimal thermostability, and stability to active detergents, builders and the like. In this respect bacterial or fungal enzymes are preferred, such as bacterial amylases and proteases, and fungal cellulases.

"Detersive enzyme", as used herein, means any enzyme having a cleaning, stain removing or otherwise beneficial effect in a laundry detergent composition. Preferred enzymes for laundry purposes include, but are not limited to, proteases, cellulases, lipases, amylases and peroxidases.

Enzymes are normally incorporated into detergent compositions at levels sufficient to provide a "cleaning-effective amount". The term "cleaning-effective amount" refers to any amount capable of producing a cleaning, stain removal, soil removal, whitening, deodorizing, or freshness improving effect on substrates such as fabrics. In practical terms for current commercial preparations, typical amounts are up to about 5 mg by weight, more typically 0.01 mg to 3 mg, of active enzyme per gram of the detergent composition. Stated otherwise, the compositions herein will typically comprise from 0.001% to 5%, preferably 0.01%-1% by weight of a commercial enzyme preparation. Protease enzymes are usually present in such commercial preparations at levels sufficient to provide from 0.005 to 0.1 Anson units (AU) of activity per gram of composition. Higher active levels may be desirable in highly concentrated detergent formulations.

Suitable examples of proteases are the subtilisins which are obtained from particular strains of *B. subtilis* and *B. licheniformis*. One suitable protease is obtained from a strain of *Bacillus*, having maximum activity throughout the pH range of 8-12, developed and sold as ESPERASE® by Novo Industries A/S of Denmark, hereinafter "Novo". The preparation of this enzyme and analogous enzymes is described in GB 1,243,784 to Novo. Other suitable proteases include ALCALASE® and SAVINASE® from Novo and MAXATASE® from International Bio-Synthetics, Inc., The Netherlands; as well as Protease A as disclosed in EP 130,756 A, January 9, 1985 and Protease B as disclosed in EP 303,761 A, April 28, 1987 and EP 130,756 A, January 9, 1985. See also a high pH protease from *Bacillus* sp. NCIMB 40338 described in WO 9318140 A to Novo. Enzymatic detergents comprising protease, one or more other enzymes, and a reversible protease inhibitor are described in WO 9203529 A to Novo. Other preferred proteases include those of WO 9510591 A to Procter & Gamble. When desired, a protease having decreased adsorption and increased hydrolysis is available as described in WO 9507791 to Procter &

Gamble. A recombinant trypsin-like protease for detergents suitable herein is described in WO 9425583 to Novo.

Cellulases usable herein include both bacterial and fungal types, preferably having a pH optimum between 5 and 10. U.S. 4,435,307, Barbesgaard et al., March 6, 1984, discloses suitable fungal cellulases from *Humicola insolens* or *Humicola* strain DSM1800 or a cellulase 212-producing fungus belonging to the genus *Aeromonas*, and cellulase extracted from the hepatopancreas of a marine mollusk, *Dolabella Auricula Solander*. Suitable cellulases are also disclosed in GB-A-2.075.028; GB-A-2.095.275 and DE-OS-2.247.832. CAREZYME® and CELLUZYME® (Novo) are especially useful. See also WO 9117243 to Novo.

Suitable lipase enzymes for detergent usage include those produced by microorganisms of the *Pseudomonas* group, such as *Pseudomonas stutzeri* ATCC 19.154, as disclosed in GB 1,372,034. See also, the lipase in Japanese Patent Application 53,20487, laid open Feb. 24, 1978. This lipase is available from Amano Pharmaceutical Co. Ltd., Nagoya, Japan, under the trade name Lipase P "Amano," or "Amano-P." Other suitable commercial lipases include Amano-CES, lipases ex *Chromobacter viscosum*, e.g. *Chromobacter viscosum* var. *lipolyticum* NRRLB 3673 from Toyo Jozo Co., Tagata, Japan; *Chromobacter viscosum* lipases from U.S. Biochemical Corp., U.S.A. and Disoynt Co., The Netherlands, and lipases ex *Pseudomonas gladioli*. LIPOLASE® enzyme derived from *Humicola lanuginosa* and commercially available from Novo, see also EP 341,947, is a preferred lipase for use herein.

The enzyme-containing compositions herein may optionally also comprise from about 0.001% to about 10%, preferably from about 0.005% to about 8%, most preferably from about 0.01% to about 6%, by weight of an enzyme stabilizing system. The enzyme stabilizing system can be any stabilizing system which is compatible with the detergent enzyme. Such a system may be inherently provided by other formulation actives, or be added separately, e.g., by the formulator or by a manufacturer of detergent-ready enzymes. Such stabilizing systems can, for example, comprise calcium ion, boric acid, propylene glycol, short chain carboxylic acids, boronic acids, and mixtures thereof, and are designed to address different stabilization problems depending on the type and physical form of the detergent composition.

E) Detergent Composition Preparation

The detergent compositions according to the present invention can be in liquid, paste or granular form. Such compositions can be prepared by combining the essential and optional components in the requisite concentrations in any suitable order and by any conventional means.

Granular compositions, for example, are generally made by combining base granule ingredients, e.g., surfactants, builders, water, etc., as a slurry, and spray drying the resulting slurry

to a low level of residual moisture (5-12%). The remaining dry ingredients, e.g., granules of the essential cellulosic based fabric treatment materials, can be admixed in granular powder form with the spray dried granules in a rotary mixing drum. The liquid ingredients, e.g., solutions of the essential cellulosic based fabric treatment materials, enzymes, binders and perfumes, can be sprayed onto the resulting granules to form the finished detergent composition. Granular compositions according to the present invention can also be in "compact form", i.e. they may have a relatively higher density than conventional granular detergents, i.e. from 550 to 950 g/l. In such case, the granular detergent compositions according to the present invention will contain a lower amount of "inorganic filler salt", compared to conventional granular detergents; typical filler salts are alkaline earth metal salts of sulphates and chlorides, typically sodium sulphate; "compact" detergents typically comprise not more than 10% filler salt.

Liquid detergent compositions can be prepared by admixing the essential and optional ingredients thereof in any desired order to provide compositions containing components in the requisite concentrations. Liquid compositions according to the present invention can also be in "compact form", in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. Addition of the cellulosic based polymer or oligomer materials to liquid detergent or other aqueous compositions of this invention may be accomplished by simply mixing into the liquid solutions the desired cellulosic based fabric treatment materials.

F) Fabric Laundering Method

The present invention also provides a method for laundering fabrics in a manner which imparts fabric appearance benefits provided by the cellulosic based polymer or oligomer materials used herein. Such a method employs contacting these fabrics with an aqueous washing solution formed from an effective amount of the detergent compositions hereinbefore described or formed from the individual components of such compositions. Contacting of fabrics with washing solution will generally occur under conditions of agitation although the compositions of the present invention may also be used to form aqueous unagitated soaking solutions for fabric cleaning and treatment. As discussed above, it is preferred that the washing solution have a pH of less than about 10.0, preferably it has a pH of about 9.5 and most preferably it has a pH of about 7.5.

Agitation is preferably provided in a washing machine for good cleaning. Washing is preferably followed by drying the wet fabric in a conventional clothes dryer. An effective amount of a high density liquid or granular detergent composition in the aqueous wash solution in the washing machine is preferably from about 500 to about 7000 ppm, more preferably from about 1000 to about 3000 ppm.

G) Fabric Conditioning

The cellulosic based polymer or oligomer materials hereinbefore described as components of the laundry detergent compositions herein may also be used to treat and condition fabrics and textiles in the absence of the surfactant and builder components of the detergent composition embodiments of this invention. Thus, for example, a fabric conditioning composition comprising only the cellulosic based fabric treatment materials themselves, or comprising an aqueous solution of the cellulosic based fabric treatment materials, may be added during the rinse cycle of a conventional home laundering operation in order to impart the desired fabric appearance and integrity benefits hereinbefore described.

EXAMPLES

The following examples illustrate the compositions and methods of the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention.

The first step in the production of the cellulosic based polymer or oligomer materials of the present invention is to convert the carboxymethylcellulose ("CMC") to a methyl ester. While there are numerous methods for accomplishing this step of the process known to those skilled in the art, two exemplary processes are given below as Examples I and II. Examples I and II differ both in the method of converting the CMC to a methyl ester as well as the feedstock of CMC as detailed below.

EXAMPLE I

Conversion of Purified CMC to Partial Methyl Ester

An 80g (0.257 mole equivalents) sample of carboxymethylcellulose (Aldrich, MW = 90,000, $DS_{RC} = 0.7$, and corresponding to the general formula defined above wherein $x = 0$) is suspended in a mixture of 196 g t-butanol, 15.7 g water, and 2.36g toluene (as an internal standard) in a 500 ml, 3-necked, round bottom flask equipped with mechanical stirrer, internal thermometer, condenser, and inert gas inlet. Then, 9.72g (0.077 mol) of dimethyl sulfate is added with stirring at room temperature. After about 2 hours, 50% sodium hydroxide is added as necessary to restore the pH to about 7-8 as judged on an aliquot diluted with 5 volumes of water. Stirring is continued at room temperature for 40 hours. At this point a 1g sample of the liquor is withdrawn and 0.25g of pyridine is added. NMR spectroscopy of this sample in dimethyl sulfoxide reveals that little or no methylated pyridine has formed showing that the dimethyl sulfate has all reacted. The modified CMC is collected by filtration and washed on the filter with several volumes of t-butanol-water and then dried under vacuum at 100°C. The atr-infrared spectrum of

this material shows carbonyl peaks for both ester and carboxylate groups consistent with the desired partial methyl ester.

The procedure is repeated except that the temperature is held at 25°C for 2 hours and then raised to 70°C for 2 hours before isolation.

EXAMPLE II


Conversion of Commercial Crude CMC to Partial Methyl Ester

An 80g (0.169 mole equiv.) sample of commercial CMC (Penn-Carbose, MW = 250,000, DS_{RC} = 0.59, 70% active) is suspended in a mixture of 197g t-butanol, 15g water, and 15g toluene (as an internal standard) in a 500 ml, 3-necked round bottom flask equipped with mechanical stirrer, internal thermometer, condenser, and inert gas inlet. Then, 14.98g (0.119 mol) of dimethyl sulfate is added with stirring at room temperature. After about 2 hours, 50% sodium hydroxide is added as necessary to restore the pH to about 7-8 as judged on an aliquot diluted with 5 volumes of water. Stirring is continued at room temperature for 2 hours and then the temperature is raised to 70°C for 2 more hours. At this point a 1g sample of the liquor is withdrawn and 0.25g of pyridine is added. NMR spectroscopy of this sample in dimethyl sulfoxide reveals that little or no methylated pyridine has formed showing that the dimethyl sulfate has all reacted. The modified CMC is collected by filtration. One half of the collected solids is recombined with one half of the liquors and saved for later reaction. The other half of the collected solids is washed on the filter with several volumes of t-butanol-water and then dried under vacuum at 100°C. The atr-infrared spectrum of this material shows carbonyl peaks for both ester and carboxylate groups consistent with the desired partial methyl ester.

EXAMPLE III

Treatment of CMC Methyl Ester with Dimethylaminopropylamine

The following is one process for converting the CMC methyl esters, such as those produced in Examples I and II above, to a partial DMAPA Amide.



One half of the CMC methyl ester produced in either of Examples I or II above is resuspended in the t-butanol-water liquor in which it was made and is treated with 3-dimethylaminopropylamine ("DMAPA", 24.2g, 0.24 mol, Aldrich) by heating under reflux for 18 hours. The solids are collected on a filter and washed with ethanol. They are then suspended in ethanol containing 10% water and the pH is raised to 11.5. After stirring at room temperature for 30 minutes, the solids are collected on a filter and washed with more aqueous ethanol and then with ethanol. An aliquot of this material is hydrolyzed by heating in 3N H₂SO₄ in D₂O for 18

hours. An aliquot of the hydrolysate is adjusted to pH 12, centrifuged to clarify, and then examined by NMR spectroscopy. This shows a resonance for the dimethylamino groups of the DMAPA and the relative size of the peak indicates that at least about 5% of the carboxylate groups have been converted to amides of DMAPA.

The treatment with DMAPA is repeated using the isolated methyl ester except that the suspending medium is toluene. The partial methyl ester is again partially converted to the DMAPA amide.

EXAMPLE IV

Partial Hydrolysis of Partial Methyl Ester of CMC

The following is another process for converting the CMC methyl esters, such as those produced in Example II above, to a partial DMAPA Amide.




A commercial sample of CMC (Penn-Carbose, MW = 250,000, $\text{DS}_{\text{RC}} = 0.59$, 70% active) which is partially converted to methyl ester by the method of Example II (DS of ester = 0.18) is suspended in ethanol-water and heated with enough sodium hydroxide to hydrolyze 70% of the esters. The resulting partially hydrolyzed ester is collected on a filter and washed with ethanol and dried under vacuum at 100°C . The atr-IR shows some residual ester carbonyl peak. The partially hydrolyzed ester displays partial solubility in water.

The hydrolysis is repeated, but the base used is an excess of 3-dimethylaminopropylamine. The isolated product shows a small amount of residual ester by atr-IR.


What is claimed is:

1. A detergent composition characterized by:

- a) from 1% to 80% by weight of surfactants selected from the group consisting of nonionic, anionic, cationic, amphoteric, zwitterionic surfactants, or mixtures thereof; and
- b) from 0.1% to 5.0% by weight of a mixture of cellulosic based polymers or oligomers of the general formula:




wherein each R is independently selected from the group consisting of



wherein:

- each R_2 is independently selected from the group consisting of H and C_1 - C_4 alkyl;
- each R_3 is independently selected from the group consisting of



wherein:

- each R_4 is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl, hydroxyalkyl, Na, K, $1/2\text{Ca}$, and $1/2\text{Mg}$;
- each R_5 is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl,

alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl and hydroxyalkyl

wherein:

each x is from 0 to 5;


each y is from 1 to 5;

provided that:


if R_4 bears a positive charge, it is balanced by a suitable anion; and

the Degree of Substitution for group R_3 is between 0.2 and 2.0, more preferably between 0.3 and 1.0, and most preferably between 0.4 and 0.7.

2. The detergent composition of claim 1, wherein at least 0.1 of the R_3 groups are derived from $-\text{CH}_2-\text{COOH}$ or a salt thereof.
3. The detergent composition of claim 1, wherein the cellulosic based polymer or oligomer has an average molecular weight of from 10,000 to 2,000,000.
4. The detergent composition of claim 1, wherein the cellulosic based polymer or oligomer has an average molecular weight of from 50,000 to 1,000,000.
5. A laundry additive composition characterized by:
 - a) from 1% to 80% by weight of water; and
 - b) from 0.1% to 80.0% by weight of cellulosic based polymers or oligomers of the general formula:




wherein each R is independently selected from the group consisting of



wherein:

- each R_2 is independently selected from the group consisting of H and C_1 - C_4 alkyl;
- each R_3 is independently selected from the group consisting of



wherein:

- each R_4 is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl, hydroxyalkyl, Na, K, $1/2\text{Ca}$, and $1/2\text{Mg}$;
- each R_5 is independently selected from the group consisting H, C_1 - C_{20} alkyl, C_5 - C_7 cycloalkyl, C_7 - C_{20} alkylaryl, C_7 - C_{20} arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl and hydroxyalkyl

wherein:

each x is from 0 to 5;

each y is from 1 to 5;

provided that:

if R_4 bears a positive charge, it is balanced by a suitable anion; and


the Degree of Substitution for group R_3 is between 0.2 and 2.0, more preferably between

0.3 and 1.0, and most preferably between 0.4 and 0.7.


6. The laundry additive composition of claim 5, wherein the cellulosic based polymer or oligomer has an average molecular weight of from 10,000 to 2,000,000.

7. The laundry additive composition of claim 5, wherein the cellulosic based polymer or oligomer has an average molecular weight of from 50,000 to 1,000,000.

8. A cellulosic based polymer or oligomer material produced by converting a carboxylalkylated cellulosic at least partially to an ester, followed by treating the converted carboxylalkylated cellulosic material with an amine in the presence of at least traces of water, wherein the cellulosic based polymer or oligomer material produced is of the general formula:




wherein each R is independently selected from the group consisting of



wherein:

- each R₂ is independently selected from the group consisting of H and C₁-C₄ alkyl;
- each R₃ is independently selected from the group consisting of



wherein:

- each R₄ is independently selected from the group consisting H, C₁-C₂₀ alkyl, C₅-C₇ cycloalkyl, C₇-C₂₀ alkylaryl, C₇-C₂₀ arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl, hydroxyalkyl, Na, K, 1/2Ca, and 1/2Mg;
- each R₅ is independently selected from the group consisting H, C₁-C₂₀ alkyl, C₅-C₇ cycloalkyl, C₇-C₂₀ alkylaryl, C₇-C₂₀ arylalkyl, substituted alkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, piperidinoalkyl, morpholinoalkyl, cycloalkylaminoalkyl and hydroxyalkyl

wherein:

each x is from 0 to 5;

each y is from 1 to 5;

provided that:

if R_4 bears a positive charge, it is balanced by a suitable anion; and

the Degree of Substitution for group R_3 is between 0.2 and 2.0, more preferably between 0.3 and 1.0, and most preferably between 0.4 and 0.7.

9. The cellulosic based polymer or oligomer material of claim 8, wherein the cellulosic based polymer or oligomer has an average molecular weight of from 10,000 to 2,000,000.

10. The cellulosic based polymer or oligomer material of claim 8, wherein the cellulosic based polymer or oligomer has an average molecular weight of from 50,000 to 1,000,000.

H:\SRCCASES\6835P\6835-epo:lp

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/19139

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C08B11/20 C08B15/00 C11D3/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C08B C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P, L	WO 98 29528 A (HUNTER KATHLEEN BRENNER ;WASHINGTON NODIE MONROE (US); WANG JIPING) 9 July 1998	1-7
A	see claims 1,2,7-9; example I; tables C,H	8-10
X	GB 2 219 587 A (GOLDSCHMIDT AG TH) 13 December 1989 see page 24, line 17 - line 21; claims 2,3	1,8
X	WO 94 24169 A (AKZO NOBEL NV ;BATELAAN JAN GERARDUS (NL); HORTS VAN DER PETER MAR) 27 October 1994	8
A	see page 11, line 12 - line 26; claims 2,3	1
A	EP 0 615 979 A (FIDIA SPA) 21 September 1994 see claims 1,7,23,32; examples 8-10,12	1,8
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 January 1999

Date of mailing of the international search report

02/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3018

Authorized officer

Loiselet-Taisne, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/19139

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 963 664 A (YALPANI MANSSUR ET AL) 16 October 1990 see claims; examples 35,36 -----	8
A	EP 0 634 481 A (PROCTER & GAMBLE) 18 January 1995 see page 2, line 27 - line 40; claims 1-8 -----	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/19139

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9829528 A	09-07-1998	NONE	
GB 2219587 A	13-12-1989	DE 3820031 A FR 2632646 A US 4988806 A	14-12-1989 15-12-1989 29-01-1991
WO 9424169 A	27-10-1994	CA 2160542 A CN 1123551 A CZ 9502658 A DE 69408454 D DE 69408454 T EP 0702697 A ES 2113646 T JP 8508538 T	27-10-1994 29-05-1996 13-03-1996 12-03-1998 20-08-1998 27-03-1996 01-05-1998 10-09-1996
EP 0615979 A	21-09-1994	AT 114322 T AU 629551 B AU 3571889 A CA 1336087 A DE 68919435 D DE 68919435 T DK 10890 A WO 8910940 A EP 0342557 A ES 2063779 T IL 90273 A JP 2504164 T US 5122598 A US 5466461 A	15-12-1994 08-10-1992 29-11-1989 27-06-1995 05-01-1995 06-07-1995 12-03-1990 16-11-1989 23-11-1989 16-01-1995 27-11-1995 29-11-1990 16-06-1992 14-11-1995
US 4963664 A	16-10-1990	US 4952684 A US 4959461 A	28-08-1990 25-09-1990
EP 0634481 A	18-01-1995	AU 7330594 A CZ 9600111 A HU 74027 A JP 9500166 T WO 9502678 A	13-02-1995 12-06-1996 28-10-1996 07-01-1997 26-01-1995